




Physics Highlights and Challenges: Astroparticle Physics

Jocelyn Monroe,
Royal Holloway, University of London

1st Joint ECFA-NuPECC-ApPEC Seminar
Fundamental Laws of the Universe in Orsay
October 14, 2019





#1: What is the universe made of?

FUTURE OF PHYSICS - THE 25 QUESTIONS



1. How did the universe begin?
2. What is the nature of the dark matter that permeates the universe?

ASTROPARTICLE PHYSICS

the European strategy



this talk:
highlights, challenges,
synergies in the context of
dark matter*




AP Associated Press

updated 5:15 p.m. ET, Sun., Aug. 12, 2007


Whoever discovers the nature of dark matter would solve one of modern science's greatest mysteries and be a shoo-in for the Nobel Prize.

UK Particle Physics Roadmap


cosmic rays




infrared/optical




X-rays




gamma-rays




radio/microwave




neutrinos



gravitational waves




dark matter?




*The tools of astroparticle physics
probe the largest and smallest distance
scales in the universe...*

The Standard Model of Cosmology




The Standard Model of Cosmology



The Standard Model of Particle Physics describes <5% of the universe!

Dark Matter Model Space: Theorist's View




(thanks to
H. Murayama)

Big Questions for the Dark Sector


(a la European Strategy for Particle Physics Update 2019)

- 1) How do we search for dark matter, depending on its properties?
- 2) What are the highlights and challenges of the experimental programs, current or near-future, that address different regions of parameter space?
- 3) How can dark matter detection experiments inform/guide energy and intensity frontier experiments, and vice versa?


Gravitational Detection




Indirect Detection



Direct Detection




Accelerator Production




Gravitational Detection




Indirect Detection



Direct Detection



Accelerator Production




Detection of Gravitational Waves: PBH?


PRIMORDIAL BLACK HOLES AS DARK MATTER

- ❖ sub-solar black holes cannot form by stellar evolution
- ❖ must be primordial in origin
- ❖ 3G detectors can probe existence of light black holes

Credit: Miguel Zumalacarregui




Future Prospects




ET in Europe, Cosmic Explorer in the US and Australia

Tension in H_0 Measurement from Standard Rulers (CMB) vs. Standard Candles (SHoES): DM Signal?




new/missing physics?




Near-future CMB polarization measurements will test new relativistic degrees of freedom + BSM neutrinos hypotheses...

Large Scale Structure Simulations




Aquarius Project, Springel+(2008)

Dark matter is the key for understanding structure formation of the universe




SIDM example that suppresses formation of low-mass subhalos:


Dark Matter
Self Interacting



Scattering off
Dark Radiation




Dissipative dark sectors can also affect small-scale structure




Observational constraints on subhalo masses are a powerful test of dark sectors, new data from e.g. GAIA, LSST will probe the CDM-predicted range


Gravitational Detection




Indirect Detection




Direct Detection




Accelerator Production




1) Self-annihilation (e.g. WIMPs)




2) Decay (e.g. sterile neutrinos)





3) Conversion (e.g. axions)



Emission energy vs. DM mass




Self-Annihilation Status and Prospects for WIMPs




Weniger, ESPPU Briefing Book

Indirect Detection Candidate Signals: Combined Fits



Testable prediction: expect anti-deuterons in AMS-02, GAPS





Korsmeier, Donato, Fornengo 2018

Leane, Slatyer 1904.08430

- We made a joint fit of the GC gamma-ray excess and of the CR excess, taking into account uncertainties in the DM distribution in the Milky Way.
- Depending on the channel there is good overlap of the two signals
- The joint-fit region is also compatible with the most recent dwarf galaxies constraints



CTA

- ⊙ New generation of instruments reaches the critical sensitivity
- ⊙ Thermal relic WIMP accessible over a very wide mass range (Galactic Centre/Halo observations @ VHE)




Decay: Sterile Neutrino Dark Matter Status, Prospects

Excess x-ray flux at 3.5 keV observed by XMM-MOS/PN, Chandra, Suzaku, NuStar in some targets but not others.




*Prospect to
test atomic line
background
hypotheses in
near-future
experiments
(XRISM ++)*


Gravitational Detection




Indirect Detection



Direct Detection




Accelerator Production



Direct Detection Strategies

kinematics: $v/c \sim 8E-4!$

recoil angle strongly correlated
with incoming WIMP direction




$$\begin{aligned} E_D &= \frac{1}{2}m_D v^2 \\ q^2 &= 2m_T E_{recoil} \\ r &= \frac{4m_D m_T}{(m_D + m_T)^2} \\ E_{recoil} &= E_D r \frac{(1 - \cos\theta)}{2} \end{aligned}$$

Spin Independent:
 χ scatters coherently off of
the entire nucleus A : $\sigma \sim A^2$
D. Z. Freedman, PRD 9, 1389 (1974)

Spin Dependent:
mainly unpaired nucleons contribute
to scattering amplitude: $\sigma \sim J(J+1)$

Observable: Recoil Energy





experimental requirements: ~1-10s of keV energy threshold, very low backgrounds

Reducible Backgrounds

Gamma ray interactions: electron recoil final states
rate $\sim N_e \times (\text{gamma flux})$, $O(1E7)$ events/(kg day)
mis-identified electrons mimic nuclear recoils

Contamination:
 ^{238}U and ^{232}Th decays,
recoiling progeny and
mis-identified alphas, betas
mimic nuclear recoils

Neutrons:
Nuclear recoil final state.
(alpha,n), U, Th fission,
cosmogenic spallation







Thanks to L. Baudis

Motivates underground laboratories, ultra-low radioactivity developments.

Irreducible Backgrounds

impossible to shield a detector from neutrino scattering!


- “neutrino floor:” both ν -N and ν -e contribute backgrounds



Modulation Signatures


Annual event rate modulation:
June-December asymmetry $\sim 2\text{-}10\%$.

Drukier, Freese, Spergel, Phys. Rev. D33:3495 (1986)



Sidereal direction modulation:
asymmetry $\sim 20\text{-}100\%$ in
forward-backward event rate.

Spergel, Phys. Rev. D36:1353 (1988)




need readout capable of direction measurement

Direct Detection Status

Wide range of parameters!

Direct detection searches historically optimised for WIMP sensitivity...





Baer et al., arXiv:1407.0017

Direct Detection Searches

Wide range of parameters


Direct detection searches



Baer et al., arXiv:1407.0017


Low Mass Prospects: Solid State Detectors

Phonons: **SuperCDMS** and **EDELWEISS** (Ge),
TES for E_{recoil} & R (timing)




Charge: biased at V , used to measure
 E_{recoil} , configured to reject surface events

detectors reach
energy thresholds
of < 1 keV, with
 $\text{FWHM} < 0.3$ keV



Phonons: **CRESST** (CaWO_4)
TES for E_{recoil}



Scintillation:
TES for particle ID





Detector Technology:
crystal bolometers with
Transition Edge Sensor
readout at $O(10 \text{ mK})$
for phonon detection +
scintillation/ionization

+ many
new ideas
(e.g. skipper
CCDs, high
pressure gas
....)



Large Mass Prospects: Liquid Noble Detectors



Detector Technology:
dual-phase
Time Projection
Chambers
with multi-tonne
liquid Xe, Ar targets

read out primary
scintillation: "S1" +
proportional gas
scintillation from
drifted electrons: "S2"

<https://lzl.slac.stanford.edu/our-research/lzl-research>

Xenon Detectors

XENON 10 (LNGS)
ZEPLIN (Boulby)

10 kg

2010



XENON 100 (LNGS)

LUX (250 kg, SURF),

PANDA-X


(500 kg, CJPL)

XMASS

(0.8t, Kamioka)

2015

Aprile et al, arXiv:1805.12565



1,000 kg

XENON 1T

(1t, LNGS)

PandaX-4:(4t, CJPL)

XENONnT: (6t, LNGS)

LZ: (7t, SURF)

2020

10,000 kg


Future: DARWIN: 50 t

Oct. 14, 2019 / p.34

Argon Detectors

DarkSide-50: leading SI limit at 1-5 GeV/c² for WIMP-nucleon and WIMP-e scattering

Agnes et al., Phys. Rev. Lett. 121.081307 (2018)



DarkSide-50
(50 kg, LNGS)




ArDM
(1t, LSC)

DEAP-3600 (3.6t, SNOLAB)



DEAP-3600:
parts-per-billion level
"S1" particle ID,
ultrapure acrylic cryostat
Ajaj et al, Phys. Rev. D (2019)



**Global Argon Dark Matter
Collaboration formed**



DarkSide-20k
(50t, LNGS)

Future: ARGO: 400 t





Jocelyn Monroe


Technology Synergies

DarkSide-20k

1. Cryostat technologies: DarkSide-20k cryostat uses technology developed for ProtoDUNE
2. Photon sensors: low noise, high efficiency, cryo Si sensors developed by DarkSide-20k with FBK (LHC Si)
3. Isotopic enhancement: ARIA facility for depletion of Ar-39 in UAr, CERN Vacuum Group collaboration



for comparison:





Aalseth, et al. JINST 12 (2017) no.09, P09030

Experiments Exploring Cryogenic SiPM Technology


Module of Opportunity for DUNE

November 12–13, 2019
Location: Brookhaven National Laboratory





Large Enriched Germanium Experiment for Neutrinoless $\beta\beta$ Decay






+ medical imaging,
environmental monitoring,
automated navigation (LIDAR) ...


Spin-Dependent Interactions

Direct Detection: leading WIMP-p limit at low mass from PICO-60, 500 kg upgrade planned. Best WIMP-n limits at high mass from Xenon-1T, at low mass from collider searches (for specific operators)

Indirect Detection: leading WIMP-p limit at high mass from neutrino telescopes, via WIMP scattering on p, +capture in the sun, leading to annihilation signatures.



Spin-Dependent Interactions: Prospects





+PINGU sensitivity
extends reach
1-2 orders of
magnitude at low
mass (<10 GeV)

+Hyper-K galactic
center search
can approach the
thermal relic bound

Direct Detection: Candidate Signal

predicted modulation $A \sim 0.02-0.1$, $t_0 = 152.5$ days

DAMA/LIBRA: measure (0.0095 ± 0.0008) cpd/kg/keV,
 $t_0 = (145 \pm 5)$ d in 2.17 tonne-yr.




CSLNGS, March 2018

many other searches, on Ge, CsI, Xe, etc.
observe no evidence of modulation.

In the same underground laboratory:
XENON100: Xe, 4.8σ exclusion of DAMA,
test of leptophilic dark matter [arXiv:1507.07748](#)



Using the same target (NaI):
ANAIS (LSC), SABRE (AU), COSINE-100 (Y2L)
~consistent at 1σ , project 3σ test in 5 years.



Direct Detection: Candidate Signal

predicted modulation $A \sim 0.02-0.1$, $t_0 = 152.5$ days

DAMA/LIBRA: measure (0.0095 ± 0.0008) cpd/kg/keV,
 $t_0 = (145 \pm 5)$ d in 2.17 tonne-yr.




CSLNGS, March 2018

many other searches, on Ge, Csl, Xe, etc.
observe no evidence of modulation.

In the same underground laboratory:
XENON100: Xe, 4.8σ exclusion of DAMA,
test of leptophilic dark matter *arXiv:1507.07748*

Using the same target (NaI):
ANAIS (LSC), SABRE (AU), COSINE-100 (Y2L)
 $\sim$ consistent at 1σ , project 3σ test in 5 years.




Direct Detection Status


Wide range of parameters!

Direct detection searches historically optimised for WIMP sensitivity...

+big recent expansion of interest in laboratory axion searches!




Baer et al., arXiv:140



Bump Hunts in Direct Detection

search for axio-electric effect:





Signal: peak in electron recoil spectrum at axion-like particle (ALP) mass.

Backgrounds: electron recoils, $\sim 1\text{E-4}/(\text{keV kg day})$.

Constraints on new pseudoscalars at $\sim 10\text{ keV}/c^2$ via ALP-electron coupling. Projected reach of DARWIN is $>\times 20$.

Constraints on vector particles at $0.1\text{-}100\text{ MeV}/c^2$ via limits on kinetic mixing to hidden sector.
(arXiv:1901.10478)

Constraints on new scalar (and vector) bosonic SuperWIMPs in $10\text{-}100\text{ keV}/c^2$ (arXiv:1709.02222)



PANDAX-II, arXiv:1707.07921

Axion/ALP searches: Mature Key Techniques




Helioscopes

- Build on success of CAST hosted by CERN
- Proposed BabyIAXO, leads to IAXO, with large discovery potential


Haloscopes

- ADMX (US) is leading the field
- In Europe, MadMax is new key player
- Smaller efforts developing new techniques



Light-shining-through-walls


- ALPS II is well underway
- STAX is a new idea RF based
- JURA is long term plan




Searches relevant for both QCD Axions and more general Axion-like particles (ALPs)

+many interesting new ideas in quantum measurement techniques, atom interferometers, ...


Gravitational Detection




Indirect Detection



Direct Detection



Accelerator Production




Collider Searches, Complementarity

- Limits on BR can be translated to limits in the DM-nucleon plane



$$\sigma_{\chi N} = \Gamma_{inv} \frac{8m_N^4 f_N^2}{v^2 \beta m_h^3 (m_\chi + m_N)^2} g_\chi \left(\frac{m_h}{m_\chi} \right), \quad (15) \quad \text{arXiv:1708.02245}$$

where $g_S(x) = 1$,

$g_F(x) = 2/(x^2 - 4)$, $\beta = \sqrt{1 - 4m_\chi^2/m_h^2}$, $v = 246$ GeV



Caveat: EFT validity
in Higgs-DM
interaction not
guaranteed beyond
HL-LHC





Caterina Doglioni - 2019/05/13 - European Strategy Update

24

Fixed-Target Searches

Renaissance of the fixed-target!

New sociology of DM searches.



Many new experiments planned or proposed for hidden sector searches.

Major synergistic topic of Dark Sectors group at ESPPU.

Ellis et al., ESPPU Physics Briefing Book, CERN-ESU-004 (2019)


Big Questions for the Dark Sector

(a la European Strategy for Particle Physics Update 2019)

- 1) How do we search for dark matter, depending on its properties?
- 2) What are the highlights and challenges of the experimental programs, current or near-future, that address different regions of parameter space?
- 3) How can dark matter detection experiments inform/guide energy and **intensity** frontier experiments, and vice versa?

Dark Matter Direct Neutrino Detection

Signal: $\nu N \rightarrow \nu N$ or $\nu e^- \rightarrow \nu e^-$



Backgrounds:

$$\gamma e^- \rightarrow \gamma e^-$$

$$n N \rightarrow n N$$

$$N \rightarrow N' + \alpha, e^-$$

$$\chi N \rightarrow \chi N?$$

*very similar requirements!
(and ideally also measure direction)*




What can future dark matter direct detection experiments tell us about the neutrino?

Open Questions

What is absolute mass scale?

- Details of neutrino oscillation?
 - Is $\theta_{23} = 45^\circ$?
 - CP violation?
- Mass hierarchy?
- Why is m_ν so small?
- Majorana or Dirac?
- Are there sterile neutrinos?



flavour key:

ν_e ν_μ ν_τ




What can future dark matter direct detection experiments tell us about the neutrino?

Open Questions

What is absolute mass scale?

- Details of neutrino oscillation?
 - Is $\theta_{23} = 45^\circ$?
 - CP violation?
- Mass hierarchy?
- Why is m_ν so small?
- Majorana or Dirac?
- Are there sterile neutrinos?

(Xe)
(Ar)



flavour key:


ν_e ν_μ ν_τ

ν -less Double Beta Decay


The liquid Xenon dark matter searches aim for competitive sensitivity to neutrinoless double beta decay.

example:
projected
sensitivity
in LZ:

Q-value=
2458 keV



P. Bras, IDPASC 2018




big opportunity:
significant Xe-136
target mass (~600 kg)

big challenges:
backgrounds,
energy resolution,
and nuclear
matrix element
uncertainty


ν -less Double Beta Decay

The liquid Xenon dark matter searches aim for competitive sensitivity to neutrinoless double beta decay.



example:
Xenon1T
energy
resolution

Q-value=
2458 keV




big opportunity:
significant Xe-136
target mass (~600 kg)



big challenges:
backgrounds,
energy resolution,
and nuclear
matrix element
uncertainty

recent demonstration of sensitivity to *rare* processes: Xe-124
two-neutrino double e- capture, *arXiv:1904.11002*




What ν physics can future dark matter detectors do?



What ν physics can future dark matter detectors do?




What ν physics can future dark matter detectors do?



Is the Neutrino Bound the End? No.

- sensitivity scales with $\sqrt{\text{time}}$ instead of linearly in time (with zero background)




(energy, angle, time) of neutrino background vs. DM signal differ.

- no ν bound for directional detectors
Grothaus, Fairbairn, JM, Phys.Rev.D90 (2014)

A ν background paradigm...
for non-directional detectors

the discovery reach
depends on ν flux errors
and on ν -e discrimination.



Conclusions & Outlook

Astroparticle experiments show that the Standard Model is incomplete!

Dark matter searches have synergies across ApPEC, ECFA, NuPPEC science.

Dark matter searches drive rapid detector technology innovation, which supports detector R&D and technology development across fields.

Astroparticle physics probes above the center of mass energy of the LHC, perhaps can tell us the next energy scale for new physics!

Astroparticle experiments continue to have outstanding discovery potential.

Many new ideas for non-standard searches in new kinds of experiments
... and today's background may be tomorrow's signal. (*T. Kajita, 2015*)





More Slides

Solar ν -Electron Scattering

Via neutrino-electron elastic scattering, LAr dark matter experiments can observe the unmeasured CNO solar neutrino flux! (via spectral deformation)
+with O(500 t-y), study the “solar metallicity problem”.

Franco et al., JCAP 1608 (2016) 08
Cerdeno, Davis, Fairbairn, Vincent, JCAP 1804 (2018) 37








big opportunity:
distinguish
between
high vs. low
metallicity.

big challenges:
Rn background
suppression and
uncertainty on
cosmogenics

*Xe-136 background makes LXe CNO challenging
Baudis et al., JCAP 1401 (2014) 044,

Geo ν -Electron Scattering

PLR analysis of energy, time, and *direction* shows sensitivity at 95% CL to measure K-40 geo-neutrino flux with O(100) t-yr exposure.



example: geo-, solar-, reactor- ν -induced electron recoil directions, at LNGS.



challenge: measure the *direction* of ~ 1 MeV e^- recoils..

Leyton, Dye, JM, Nature Commun. 8 (2017) 15989


Sterile ν Signatures

The beta decay energy spectrum is modified by neutrino mass and mixing.

Upper limit on $|U_{e4}|^2$ at 10 keV mass ~ 0.02 at 90% CL from beta decay.



big opportunity:
high Q-value beta decays of backgrounds (e.g. Ar-39) with good energy resolution have sensitivity to 10-100 keV sterile ν s




big challenges:
nuclear physics uncertainties on beta spectrum shape, understanding energy resolution at % level

Complementarity: Case 0 Example

Example: Scalar DM – Scalar Mediator
 $m = 100 \text{ GeV}$

A single target cannot determine the DM mass and couplings

Xe




Complementarity: Case 0 Example

Example: Scalar DM – Scalar Mediator
 $m = 100 \text{ GeV}$

A single target cannot determine the DM mass and couplings

The experimental response is very sensitive to the target




Complementarity: Case 0 Example

Example: Scalar DM – Scalar Mediator
 $m = 100 \text{ GeV}$

A single target cannot determine the DM mass and couplings

The experimental response is very sensitive to the target



Combining data some degeneracies can be removed



Xe
 Ar

Directional Detection

R&D towards recoil *direction* measurement to correlate a signal with the galactic halo



Many R&D efforts:

DRIFT, DMTPC, MIMAC, NEWS-DM, RED, CYGNO++

largest are 1m³ (O(100g) target).


Majority use CF₄ gas; NEWS uses emulsions.

CYGNUS: global coordination towards a physics-scale directional experiment.
Physics Reports 2016, arXiv:1602.03781

huge experimental challenge to measure direction of recoil tracks of O(10 keV): <mm length!

Directional Detection

R&D towards recoil *direction* measurement to correlate a signal with the galactic halo



detectors achieve angular resolution of $\sim 35^\circ$ at 50 keVr

with current best direction reconstruction, need 200-400 events to measure anisotropy at 3σ significance

Phys.Rev.D95 (2017) 122002

Is High Mass Interesting? (1)



Yes. >few hundred GeV is above LHC reach, but accessible in direct detection experiments.

In EFT approach, the spectrum from possible interactions (e.g. momentum dependent) does not have the typical WIMP exponential.

Information isn't only at threshold!

Beyond SUSY, variety of models can have DM candidates up to few TeV, e.g.


- little Higgs,
- warped extra dimensions,
- walking technicolor
- MIMPs
- composite states
- ...



Is High Mass Interesting? (2)

Sensitivity to composite dark matter, e.g. *Hardy, Lazenby, March-Russell, West JHEP 07 (2015)*
dark nuclei, formed of k bound states of self-interacting light dark nucleons.

Scattering process now has a form factor from the nuclear dark matter and the target.



example: dark nucleon
mass = 1 GeV, $r = 1$ fm,
and per-SM nucleon
 $x_{\text{sec}} = 1\text{E-}46$ cm².

*Kirk, Butcher, JM, West,
JCAP 1710 (2017) 10, 035*

