





Search for Dark Matter with XENON and DARWIN

Julien Masbou
Subatech – Université de Nantes

WIMP direct detection principle

Nuclear
Recoil
(NR)



Electronic
Recoil
(ER)

γ and β particles
interact with the atomic electrons
→ Background in « standard
WIMP analysis »

XENON Collaboration



 ~ 170 scientists

 27 institutions

 11 countries



Julien Masbou, CosPT 2021, Orsay, 11th October 2021

Phases of the XENON Program



XENON10

2005 – 2007

15 cm drift TPC

Total: 25 kg

Target: **14** kg

ER background :
1000 evt/(t.keV.d)

Achieved (2007)

$$\sigma_{\text{SI}} = 4.5 \cdot 10^{-44} \text{ cm}^2$$

@ 30 GeV/c²



XENON100

2008 – 2016

30 cm drift TPC

Total: 161 kg


Target: **62** kg

ER background :
5.3 evt/(t.keV.d)

Achieved (2016)

$$\sigma_{\text{SI}} = 1.1 \cdot 10^{-45} \text{ cm}^2$$

@ 55 GeV/c²



XENON1T

2011 – 2018

100 cm drift TPC

Total: 3 200 kg

Target: **2 000** kg

ER background :
0.2 evt/(t.keV.d)

Achieved (2018)

$$\sigma_{\text{SI}} = 4.1 \cdot 10^{-47} \text{ cm}^2$$

@ 30 GeV/c²



XENONnT

2019 – 2025

150 cm drift TPC

Total: 8 600 kg

Target: **5 900** kg

ER background :
 $2.2 \cdot 10^{-3}$ evt/(t.keV.d)


Projected

$$\sigma_{\text{SI}} = 1.6 \cdot 10^{-48} \text{ cm}^2$$

@ 50 GeV/c²

Dual phase TPC: principle

TPC = Time Projection Chamber



S1:

→ Photon ($\lambda = 178 \text{ nm}$)
from Scintillation process


S2:

→ Electrons drift
→ Extraction in gaseous phase
→ Proportional scintillation light


3D reconstruction :

→ X,Y from top array
→ Z from Drift time


$$(S2/S1)_{\text{WIMP},n} < (S2/S1)_{\gamma,\beta}$$




Origins of backgrounds



Uses of S1 & S2 with XENON1T data




XENON1T Results – NR Searches




Standard Analysis (S1-S2)

- high mass WIMP search ($> 6 \text{ GeV}/c^2$)
 - minimum of $4.1 \times 10^{-47} \text{ cm}^2$ at $30 \text{ GeV}/c^2$ (90% C.L.)
- PRL 121 (2018) 111302



XENON1T Results – NR Searches




Standard Analysis (S1-S2)

- high mass WIMP search ($> 6 \text{ GeV}/c^2$)
 - minimum of $4.1 \times 10^{-47} \text{ cm}^2$ at $30 \text{ GeV}/c^2$ (90% C.L.)
- PRL 121 (2018) 111302

S2-only NR

- using charge signal (S2) only
 - lowering threshold
 - low-mass WIMP search ($< 6 \text{ GeV}/c^2$)
- PRL 123 (2019) 251801

XENON1T Results – NR Searches



Standard Analysis (S1-S2)

- high mass WIMP search ($> 6 \text{ GeV}/c^2$)
 - minimum of $4.1 \times 10^{-47} \text{ cm}^2$ at $30 \text{ GeV}/c^2$ (90% C.L.)
- PRL 121 (2018) 111302


S2-only NR

- using charge signal (S2) only
 - lowering threshold
 - low-mass WIMP search ($< 6 \text{ GeV}/c^2$)
- PRL 123 (2019) 251801

S2-only Migdal (NR + ER)

- using charge signal (S2) only for NR+ER
 - Migdal effect (theoretical prediction)
 - low-mass WIMP sensitivity down to $85 \text{ MeV}/c^2$
- PRL 123 (2019) 241803

XENON1T Results – NR Searches



CE ν NS Search

- Coherent Elastic Neutrino-Nucleus Scattering for ^8B neutrinos from the Sun
- irreducible background in future WIMP detectors
- we lowered our threshold from 3-fold to 2-fold PMT coincidence





Results

- no excess observed
 - use to constrain light and charge yield of detector at lowest energies
 - place new limit on light dark matter between 3 – 11 GeV/c^2
- PRL 126, 091301 (2021)

XENON1T Results – ER Searches

Abundance:



$^{124}\text{Xe} \sim 1 \text{ kg / t}$



Detectable cascade of X-rays and Auger electrons in the **keV-range (64.3 keV)**

$$T_{1/2} = (1.8 \pm 0.5_{\text{stat}} \pm 0.1_{\text{sys}}) \times 10^{22} \text{ y}$$

XENON1T Results – ER Searches



76 ± 2 events/(t·y·keV)

Fitted background in [1, 30] keV

Events in [1, 7] keV **232 ± 15** **285**
 expected observed

XENON1T Results – ER Searches



3.2 σ TRITIUM BACKGROUND

- Fitted concentration: $(6.2 \pm 2.0) \times 10^{-25}$ mol/mol $^3\text{H}/\text{Xe}$
- We don't expect that much ^3H from liquid purity
- Very difficult to confirm or exclude such a tiny abundance



3.4 σ SOLAR AXIONS

- Non-null coupling to electrons $\rightarrow$ ABC and/or Primakoff
- Strong tension with astrophysical constraints
- Axions+ ^3H favoured over ^3H -only at 2.1 σ




3.2 σ NEUTRINO MAGNETIC MOMENT μ_ν

- $\mu_\nu = [1.4, 2.9] \times 10^{-11} \mu_B$
- $\mu_\nu > 10^{-15}$ would imply neutrinos to be Majorana fermions
- Tension with astrophysical constraints





3.0 σ BOSONIC DARK MATTER

- Including pseudo-scalar (ALPS) and vector (dark photons) bosons
- Most restrictive constraints to date set




From XENON1T to XENONnT



From XENON1T to XENONnT

Larger TPC



- Total 8.6 t LXe
- 5.9 t in TPC
- ~ 4 t fiducial
- 248 $\rightarrow$ 494 PMTs



From XENON1T to XENONnT

Neutron Veto



- Inner region of existing muon veto
- optically separated
- 120 additional PMTs
- Gd in the water tank
- 0.5% $\text{Gd}_2(\text{SO}_4)_3$



From XENON1T to XENONnT

LXe purification


- Faster xenon cleaning
- 5L/min LXe (2500 slpm)
(XENON1T: 120 slpm)



From XENON1T to XENONnT

^{222}Rn Distillation


- Reduce radon from pipes, cables, cryogenic system
- New system, tested in XENON1T
- Background reduction 1/10



From XENON1T to XENONnT

Storage and Recovery (ReStoX-II)

- 10 tons capacity
- Storage in Solid / Liquid / Gas
- Fast recovery (1 ton / hour)





XENONnT expected sensitivity

E. Aprile et al JCAP11(2020)031

After 20 tons x yr,
the expected sensitivity is:


$$1.4 \times 10^{-48} \text{ cm}^2$$

@ 50 GeV/c² (90% CL)



Detailed investigation on the low-energy ER
excess observed in XENON1T



Preliminary study suggest that after few months of
XENONnT data the solar axion signal hypothesis could be
differentiated from 3H background at the 5 σ level



The DARWIN Project


- **Time-projection chamber:** 40 active tons of LXe, 2.6 m in diameter and height
- **Ultra-low background Goal:** deep underground, low background cryostat, neutron and muon veto

JCAP 1611 (2016) no.11, 017



The DARWIN Project


JCAP 10, 016 (2015)



Direct Dark Matter Search
expected sensitivity with 200 ton x yr
 $2.5 \times 10^{-49} \text{ cm}^2 @ 40 \text{ GeV}/c^2$

current limits

near future sensitivity
DARWIN's sensitivity




Neutrinoless Double Beta Decay Search
Optimised 5 t fiducial of ^{136}Xe

Sensitivity : $T_{1/2} = 2.4 \times 10^{27} \text{ y}$ (90 % CL)
 3σ discovery potential: $1.1 \times 10^{27} \text{ y}$
After 10 years of observations

Eur. Phys. J. C 80 (2020) 808

Evolution of LXe TPC as WiMP detectors



Evolution of LXe TPC as WiMP detectors

Fiducial mass [kg]

XENON10

5

XENON100

34

LUX

118


PandaX

306

XENON1T


1300

1000



Summary and outlook

- Xenon based Dual Phase Time Projection Chamber has proven to be the **leading technology** in the field of direct Dark Matter searches.
- XENONnT is **about to start its physics program**, and aims to improve the results of its predecessor and answer to the Low-ER excess question.



- DARWIN will be **much more than the ultimate LXe-based dark matter detector**
➔ large detector with ultra-low backgrounds, very good energy resolution, low energy threshold